

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

Intyg Certificate



Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande Biovelop International BV, PJ Amstelveen NL
Applicant (s)


(21) Patentansökningsnummer 0103328-1
Patent application number

(86) Ingivningsdatum 2001-10-04
Date of filing

(30) Prioritet begärd från 2001-02-26 SE 0100655-0

Stockholm, 2003-08-11

För Patent- och registreringsverket
For the Patent- and Registration Office


Christina Vängborg

Avgift
Fee 170:-

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassen

1

TITLE

PROCESS FOR THE EXTRACTION OF CELL WALL COMPONENTS AND LESS ACCESSIBLE PROTEINS FROM CEREAL BRANS SUBSTANTIALLY FREE OF SOLUBLE COMPOUNDS

DESCRIPTION

TECHNICAL FIELD

The present invention relates to process for the extraction of cell wall components and less accessible proteins from cereal brans substantially free of soluble compounds, the compounds thus recovered as well as their use.

INTRODUCTION

Bran is defined as the seed coat of cereal grains such as wheat, barley, rye, triticale, oat or rice. Anatomically, bran comprises the outer layers of the seed, known as the pericarp-testa and an inner layer known as the aleurone layer, which is often classified as the outermost layer of the endosperm. However, from the practical point of view cereal bran is herein defined as the remaining material after the conventional milling or polishing of cereal grains and contains primarily pericarp-testa and aleurone layer components, along with the cereal germ and residual parts of the endosperm. The relative amounts of each component will depend upon the type of cereal and milling technique applied.

Within this definition, bran therefore contains all of the pericarp- testa components, the aleurone layer, the germ components including germ proteins and oils, along with a residual amount of endosperm starch, gluten and pentosans.

In this patent, the term cereal bran substantially free of soluble compounds or "cleaned bran" refers to any cereal bran, which has been processed, after conventional milling or polishing, by any means so as to remove substantial amounts of soluble components, which are extracted by water or organic solvents. The resulting material, hereafter referred to as cleaned bran, should contain rather limited amounts of soluble sugars, starch and gluten (less than 1%), but it may still contain some proteins and fats, which are less accessible

Ink. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassar

2

and/or soluble. The cleaned bran consists primarily of cell wall components, of which hemicellulose is the most abundant.

5 The invention relates to methods, procedures and an industrial process for the wet-fractionation of bran into two protein rich fractions, one of which contains the germ oils and related components, a fibre fraction, which also retains most of the aleurone proteins, and a sugar syrup fraction.

10 The invention is centred around the wet-milling of bran in the presence of enzymes: a) starch degrading enzymes (polysaccharidases) such as alpha amylases, and amyloglucosidases, and b) non-starch degrading enzymes (polysaccharidases) and c) a phytase, under appropriate conditions of temperature, i.e. from 50 to 90, more preferably from 50 to 75, and pH from 4 to 7.5. This is followed by the separation of the above listed
 15 components from aqueous suspension using mainly centrifugal separation methods. The pH when using an alpha amylase is normally around 7, and when using an amyloglucosidase it is around 4.5. The enzymes are used normally in a cocktail comprising 200 to 1500 IU/g of substrate.

20 Presently there exists no commercial method for wet-fractionating cereal bran to produce the specific products mentioned above.

This invention relates to methods, procedures and an industrial process for the wet-fractionation of cleaned bran into one protein rich fraction, which contains proteins from the
 25 aleurone cells, a soluble hemicellulose fraction, a soluble oligosaccharide fraction and an insoluble fibre fraction.

This invention further aims at the fractionation of cleaned bran by combining wet-milling and enzymatic hydrolysis specifically with food grade xylanases under well-controlled
 30 conditions of temperature, i.e. from 35 to 80°C, more preferably from 40 to 50°C, and pH from 4 to 7, preferably 4.5 to 5.5.

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassen

3

This step is followed by the separation of insoluble fibre and protein fractions from aqueous suspension using centrifugal separation methods while both hemicellulose and oligosaccharide fractions are separated by size exclusion techniques such as ultrafiltration.

5

Presently there exist no commercial enzyme-based methods for wet-fractionating of cereal brans or derived products such as cleaned bran, which is capable of extracting high amounts of less accessible components, such as hemicellulose and aleurone proteins.

10 PRIOR ART

15

US patent 4,361,651 describes a process for making fermentable sugars and high protein products from grain, mainly maize. In this method, grain is steeped for 10-30 hours, prior to milling and separation of the germ component, saccharification of carbohydrates (mainly starch), and separation of fibre. The yield of starch is maximised for fermentation to alcohol. Within the described process there is no specific fractionation of the bran component, separation of protein types or consideration of the germ component.

20

US patent 5,312,636 informs on a process for fractionating crop into industrial raw material. This is focused on oat grain and incorporates bran fractionation procedures that involve the extraction of more hydrophobic components such as lipids in polar organic solvents prior to the alkaline extraction of residual bran to produce beta-glucan, protein and degummed fibres. The use of the organic solvent is a key step in the process and hydrolysing enzymes are not utilised during the fractionation procedure.

25

30

Two related US patents (4,171,383 and 4,171,384) inform on dry and wet milling procedures for refining whole wheat grain. US-A-4,171,383 focuses on wet milling of the whole kernel. The bran produced is mixed with a separated (mainly) endosperm protein fraction to produce animal feed. 4,171,383 describes dry milling of the whole kernel to produce an endosperm fraction, a germ fraction and a bran fraction. The endosperm fraction is then subjected to wet milling and separation of starch-rich and protein-rich fractions. The protein rich fraction is added to the bran to produce an animal feed. There is no description of a specific wet fractionation of the bran itself within either patent.

Ink. t. Patent- och reg.verket

2001-10- U 4

Muvudfaxen Kassen

4

WO 99/11672 patent application discloses a process that uses selective enzymes, such as acetyl xylan esterase and ferulic acid esterase, to both facilitate the removal of hemicellu-

5

lose from various plant materials and alter its degree of phenolic ester substitution. Despite the fact that functional hemicellulose with high solubility and gelling strength can be produced yields are rather low. In fact, the inventors reported a 3 and 6% yield of arabinoxylan ferulate (hemicellulose), when wheat bran was treated with acetyl xylan esterase for 90 and 180 min, respectively. Furthermore, the invention does not make any reference to the use of xylanases, or its combination with wet milling in order to overcome the low yields reported.

10

US-A- 5,308,618 discloses a process to extract soluble dietary fibre hemicelluloses from wheat bran by applying a heat pre-treatment in aqueous solution. This is followed by further processing such as filtration, salting out, dialysis, ultrafiltration, reverse osmosis, gel filtration and precipitation in order to remove contaminants from the hemicellulose fraction. The inventors make no claims with regards to the use of enzymes and production of products streams other than hemicellulose. Furthermore, the invention highlights the need of run costly procedures to remove contaminants, which were once present in the original wheat bran. The bran is extracted at high temperatures and pressures in water (180 - 200° C), producing a glucose rich dietary fibre component in the water phase. The process specifically targets the production of dietary fibre and is not really / strictly a fractionation procedure in that other products are largely ignored.

15

20

25

US-A- 3,879,373 discloses a process to extract hemicelluloses from wheat bran by applying alkali treatment to dissolve hemicelluloses and other bran components followed by ethanol extraction to separate the hemicelluloses. Alkali (sodium hydroxide) extraction of hemicellulose has also been disclosed in US-A- 5,174,998 as an intermediate step to produce controlled-release compositions containing the said alkali-extracted hemicellulose and an active substance. Similar alkali-extraction procedure is disclosed in US-A- 4,927,649 to produce hemicellulose, which is then used in coating compositions containing insoluble dietary fibre.

30

Ink. t. Patent- och reg.verket

2001-10- U 4

Muvudfaxen Kassen

5

WO 00/04053 patent application describes a chemical process using alkaline peroxide treatment to produce high yields of light coloured gelling hemicelluloses from products derived from flour, husk or bran. Another chemical extraction process of hemicellulose from wheat bran has been disclosed in WO98/31713 patent application, whereby the inventors combine a washing procedure to remove the starch fraction followed by an alkaline treatment with sodium hydroxide to extract the hemicellulose from the starch-free raw material.

10 It appears from above-described prior arts on alkali extraction of hemicellulose that this is an old, proven and effective way to yield high quantities of soluble hemicellulose with interesting functionalities such as gelling, dietary fibre and as an inert material for controlled-release compositions. The drawback of such technology is the associated problems of utilising chemicals. Firstly, chemicals eventually become contaminants in various product streams, and therefore require additional purification. This normally has significant cost implications. Secondly, innovative industrial processes based on chemical extraction are not always attractive from the marketing point of view, particularly in food applications.

20 Production of insoluble dietary fibres from oats is disclosed in US-A- 5,023,103, which describes a chemical procedure (alkali and bleaching treatment) for the production of insoluble dietary fibre with high water holding capacity and non-gritty mouth feel. A water holding capacity of 6.9g water/g oat fibre has been reported.

25 Other references have disclosed interesting processes for the extraction of proteins from cereal brans. US-A- 4,746,073 discloses a physical process to separate aleurone cell particles and pericarp-testa particles from commercial wheat bran. The process consists of milling the bran particles to a specific particle size distribution, electrostatically charge the said particles and then pass the said charged particles through a magnetic field, which separates aleurone from pericarp-testa particles. This is a rather different concept from the current invention, which is based upon the use of enzymes and wet milling. The separation is achieved by hammer-milling the bran and then subjecting the resultant particles to a

Ink. t. Patent- och reg.verket

2001-10- U 4

6

Huvudfaxen Kassar

physical separation regime. No wet processing is employed during the fractionation procedure described therein.

5 Waszczynskyj et al. (1981) have proven that protein extraction rate of alkali-treated full fat wheat bran can be increased from 30% up to 38.5% when it is preceded by polysaccharidase treatment. The above-mentioned figures are significantly lower than those described in the present invention whereby up to 60% protein extraction rates were achieved without using alkali treatment. Furthermore, US 5,622,738 discloses a method to extract soluble
 10 hemicelluloses, for use as a source of dietary fibre, from various fibrous materials including cereal brans using alkali digestion followed by xylanase treatment. As in other prior arts, the inventors made use of alkali digestion to improve extraction rates. Additionally, the residence time for the enzymatic treatment was rather long (3 to 96h), which makes the process not very attractive from the production cost point of view.

15

It is clear that none of the above-mentioned prior arts have succeeded to simultaneously yield aleurone proteins, oligosaccharides and hemicelluloses, and yet produce insoluble dietary fibre from previously cleaned cereal bran, i.e. substantially free of soluble components, using xylanases in combination with wet milling.

20

Furthermore, the current invention has arrived at an industrial process, which enables one to extract high amounts of such components (up to 50%) without using alkali digestion.

The main objectives of this invention were to:

25

1. Arrive at an efficient and cost effective industrial wet process to extract and yield hemicellulose, oligosaccharides, aleurone proteins and insoluble dietary fibre from cleaned cereal brans.

2. Combine the use of xylanases treatment with wet milling to improve the efficiency of extraction and separation in an industrial process.

30

3. Ensure that the raw material contains the least amount of readily extractable components, hence solubles, so that contamination with the said solubles in the end products is kept to a minimum.

Ink. t. Patent- och reg.verket

2001-10- U 4

Muvudfaxen Kassen

7

4. The process is carried out preferably with food grade and non-genetically modified (non-GMO) xylanases in order to broaden the market opportunities for the end products.

5 DESCRIPTION OF THE PREFERRED EMBODIMENTS

Various methods of extraction and fractionation of hemicelluloses from cereal brans have been developed in the past. Equally, various methods for the extraction of valuable proteins and insoluble dietary fibre from cereal brans have been disclosed. The problem is that when one combines the use of commercial cereal brans, which contain large quantities of soluble components such as starch, soluble proteins, pentosans, oils, etc, and a simple extraction process such solubles eventually become contaminants of the main product streams, and therefore have to be removed. This is a costly procedure and in many cases jeopardizes the market value of the non-hemicellulose components.

15

EXAMPLE 1

Wheat bran produced from short milling (SMB) and conventional milling (CMB) processes were used in this trial. Bran sample of 25 kg was transferred to a mixing tank and sequentially hydrolysed at temperatures varying from 70 °C at the first stage with α -amylase to 60 °C in the second stage with amyloglucosidase for a total hydrolysis time of 4 h. During this period the reaction mixture was intermittently wet milled to increase in surface area and dispersion of soluble components. The pH of the reaction mixture was set at neutral initially and then decreased down to 4.5 with acetic acid in the second stage. In addition of maximising the enzymatic activity the acidic pH allowed partial solubilization of the phytates present in the bran.

25

At the end of the enzymatic hydrolysis - wet milling step the enzymes contained in the reaction mixture were inactivated by wet heating through a heat exchange and quickly cooled down to room temperature.

30

The hydrolysed bran solution was then put through a two-phase decanter to separate the insoluble (fibre and aleurone fractions) from the soluble fraction.

Ink. t. Patent- och reg.verket

2001-10- U 4

Muvudfaxen Kassar

8

The soluble fraction was put through a separator so that the heavy phase containing mostly the germ components could be separated from the light phase containing mostly components from the remaining endosperm found in the bran. The light fraction, which was heavily contaminated with sugars, was put through an ultrafilter having a 50 kDa filter in order to separate low molecular weight sugars and a protein fraction with less sugar contamination from each other.

All soluble protein fractions, i.e. heavy and light phases, were blended together and finally processed through spray drying. The sugar fraction was concentrated by vacuum evaporation at mild temperature (40 to 60 °C) until a 75% sugar concentration was achieved. The fibre fraction was dried in a conventional laboratory oven, but in an industrial process this can be carried out by a number of different dryers, i.e. tumble drier, ring drier, fine grinder, etc.

Average chemical composition of the brans and their respective fractions are shown below in Table 1.

TABLE 1

Sample	Dry matter	Protein	Oil	Fibre	Ash	NNE***
CMB*	90.8	15.7	4.1	45.4	5.5	29.3
CMB fractions of the process						
Protein phase	92.9	31.8	7.7	1.1	7.9	51.5
Fibre	92.8	13.6	3.0	76.9	4.1	2.4
SMB**	89.1	14.3	2.3	23.7	3.2	56.5
SMB fractions of the process						
Protein phase	93.9	27.8	1.5	0.9	3.4	66.4
Fibre	94.3	22.5	4.1	64.8	1.6	7.0

* Conventional milling bran

** Short milling bran

*** Non-nitrogen extracts

Int. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassen

9

EXAMPLE 2

Wheat bran produced from conventional milling was subjected to enzymatic treatment and wet milling as described in Example 1. The hydrolysed bran was fractionated using a two-phase decanter into an insoluble (combined fibre and aleurone) and a soluble fraction.

5

The soluble fraction was fed into a separator for fractionation using centrifugal forces thus producing two phases. The germ-rich phase was washed with water and fed again to the separator to remove the excess sugars and other light contaminants. The resulting protein fraction was kept as such or mixed with evaporated liquid whey on a 1:1 ratio (dry matter basis).

10

The endosperm-rich wheat fraction, which was heavily contaminated with sugars, was fed to an ultrafilter in order to separate low molecular weight sugars and a protein fraction with less sugar contamination.

15

All soluble protein fractions, i.e. germ and endosperm-rich phases and the mixtures with whey, were spray dried separately. The sugar fraction was concentrated by vacuum evaporation at mild temperature ($t = 60\text{ }^{\circ}\text{C}$) until a 75% sugar concentration was achieved. The fibre fraction was oven dried.

20

The additional washing carried out on both germ-rich protein and fibre fraction was very effective to decrease the amount of light soluble contaminants from each fraction, and therefore increase the relative content of valuable components.

25

Compositional data indicates that germ and endosperm-rich protein fractions have a different relative content of protein and oil. Protein and oil content from the former were 48.6 and 18.6%, respectively and those from the latter were 28.7% and 1.5%, respectively. The insoluble phase containing primarily the bran pericarp (fibre) and the aleurone proteins had 86.4% fibre and 12.6% protein. The chemical composition of the germ-rich phase -

30

whey mix was 31.5% protein, 9.8% oil and 37% lactose.

A further important observation was that the spray dried germ-rich fraction containing 18.6% oil was substantially more resistant to oxidation (rancidification) compared to the

Ink. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassen

10

original wheat bran. The original wheat bran started getting rancid after 3 weeks of storage. Despite the fact that no exogenous anti-oxidants were added to the germ-rich fraction it only started going off after 12 weeks of storage.

5

EXAMPLE 3

Previous examples illustrate the use of starch-hydrolysing enzymes and wet milling followed by various separation steps in order to yield both protein, sugar and fibre fractions, the latter still containing substantially high amounts of aleurone proteins. It could be of interest for some applications to separate, at least partly, the aleurone proteins from the bran pericarp (fibre) and recover such proteins in the same fraction as the endosperm-rich fraction for instance.

10

A trial was set up in the same way as described in EXAMPLE 2, except that a cocktail of polysaccharidases containing both high cellulase and xylanase activities was added together with the amyloglucosidases, and let to work for 3 h. Temperature and pH conditions were kept unchanged. The resulting reaction mixture was further treated exactly as described in EXAMPLE 2.

15

The inclusion of polysaccharidase during the hydrolysis step had a positive effect with regards to aleurone protein extraction and protein recovery as measured by the mass balance and protein content. The protein content in the endosperm-rich fraction increased from 28.7% (without polysaccharidases) to 34.7% (with polysaccharidases) and the overall protein recovery was increased by 35% when polysaccharidase was added.

20

25

EXAMPLE 4

The colour of protein ingredients can be of importance particularly in some food and feed applications. Milk products such as caseinates, whey powder and whey protein concentrate have a light colour and soy protein concentrate have a light brown colour. These products are the main ingredients in high value feeds such as calf milk replacer. But, in some food applications such as sausage and hamburger despite the fact the inclusion level is much lower, colour can still play an important role in the product acceptability.

30

Ink. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kesson

11

The technical feasibility of bleaching the germ-rich fraction was assessed by two means. 1. Solely alkali and hydrogen peroxide bleaching, and 2. Alkali-free peroxidase and hydrogen peroxide bleaching.

5

1. 10 (ten) g samples of germ-rich fraction were incubated in 1 L beakers containing 100 ml water. Samples were dispersed with stirring and ca. 0.25ml NaOH added until pH 12 was reached. Solutions were warmed at 50 °C and 3.5, 5 and 10 ml of 30% H₂O₂ were added to different flasks to provide uptake levels 10, 15 and 30% H₂O₂ on weight basis of germ-rich
 10 fraction. Mixtures were stirred for 1 h and neutralised with acetic acid.

Full bleaching was achieved with 15 and 30% H₂O₂. Sample treated with 10% H₂O₂ was only partly bleached. All alkali bleached samples became darker with drying.

15 2. 10 (ten) g sample of germ-rich fraction was incubated in 1 L beaker containing 100 ml water. Samples were dispersed with stirring and ca. 0.25 ml NS 51004 Novozymes peroxidase was added. Solution was warmed at 50 °C and 3.5 of 30% H₂O₂ was added to the flask, i.e. 10% H₂O₂ on weight basis of germ-rich fraction, and the mixture stirred for 2 hrs.

20 The peroxidase - hydrogen peroxide bleaching was effective, consumed less chemicals and no darkening of the sample was observed after drying.

By utilising previously cleaned bran as the preferred raw material, the inventors have overcome many important production constraints and created interesting opportunities to
 25 extract and separate new components from cereal brans. Furthermore, the inventors have developed a simple method in which wet milling is combined with enzymatic treatment using food grade commercial xylanases, and cheap industrial separation processes.

Essentially, this invention allows one to economically produce proteins derived
 30 substantially from the aleurone cells, hemicellulose, oligosaccharide and insoluble dietary fibre.

Ink. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassar

12

EXAMPLE 5

Laboratory scale trials were carried out on cleaned wheat bran to test extraction rates using xylanase treatment. The cleaned wheat bran used as a raw material contained less than 1% starch and at least 50% and 70% of the protein and oil, respectively, originally found in the starting material had been removed.

Ten g cleaned bran were incubated in 150 ml water, the pH adjusted to 5.5 with acetic acid and an enzyme cocktail containing pentosanase and hemicellulase activities was added at the following concentrations: 0 (control), 0.1, 0.25, 0.5, 1 and 2% (w/w basis). Reaction mixtures were kept at 40°C for 120 min. The treatment was terminated by inactivating the enzymes at 80°C for 30 min.

Results indicated relatively high extraction rates compared to the control treatment (no enzyme added) despite the amount of enzyme used. Extraction rates of 3.1, 32.0, 32.8, 33.1, 33.8 and 34.2%, respectively, were obtained from control, 0.1, 0.25, 0.5, 1 and 2% treatments, respectively.

EXAMPLE 6

A similar trial to that described above was carried out with a purified endo 1,4-beta xylanase (pentosanase) at two levels of inclusion: 0.25 and 0.5% (w/w basis).

Extraction rates were also high, and increased from 3.1 (control treatment - no enzyme) to 28.6 and 26.1%, when 0.25 and 0.5% pentosanases were added, respectively.

EXAMPLE 7

Cleaned bran with the same specification as described in Examples 5 and 6 was used in a large-scale trial. The objective was to validate a process using standard industrial equipments, quantify process parameters, determine extraction rates of the various fractions, and ultimately characterize the end products.

Ink. t. Patent- och reg.verket

2001-10- U 4

13

Huvudfaxen Kassen

Cleaned bran (80 kg) was incubated in hydrolysis tanks containing 500 l water. The pH was adjusted to 5.5 and a purified endo 1,4-beta xylanase (pentosanase) was added at 0.5% (w/w basis). The reaction mixture was continuously stirred and intermittently wet milled while kept at 40°C for 90 min. The hydrolysis/wet milling treatment was terminated by heating up the reaction mixture to 90°C for 2 min in a heat exchange device.

The inactivated hydrolysate was pumped through a commercial two-phase decanter where the insoluble phase (insoluble dietary fibre) was separated from the solubles. The insoluble phase was dried and further milled in a commercial fine grinder using indirect heat.

The solubles were then pumped through another two-phase decanter where a heavy phase (aleurone protein-rich fraction) was separated from a light phase containing the extracted hemicellulose fraction in the form of both soluble hemicelluloses and oligosaccharides. The protein-rich phase was spray dried.

The hemicellulose fraction was further separated by size exclusion using an ultrafiltration unit whereby the large molecular size fraction (soluble hemicellulose) was separated from the small molecular size fraction (oligosaccharides and sugars). The resulting fractions were further processed by spray drying into a fine powder or alternatively evaporating the excess water until 25% water content was achieved.

The following yields of insoluble dietary fibre, hemicellulose, oligosaccharides and protein-rich fractions were obtained from cleaned wheat bran 51.0, 26.1, 17.3 and 7.7%, respectively.

EXAMPLE 8

An insoluble fibre fraction extracted according to the procedure described in Example 7 was characterized with focus on its potential use as a source of dietary fibre and texturizer in food applications.

Typical composition was as following: dry matter 95%, cell wall components 75%, protein 11%, soluble sugars 3% (of which at least 75% is glucose), fat 4% and minerals 1.5%.

Link t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassar

14

The water holding capacity (WHC), of primary importance when assessing the usefulness of insoluble dietary fibres, of the above-described product was 8.6g water/g sample on dry basis. For comparison purposes the WHC of wheat bran in the range of 3.5g/g and that of
 5 cleaned wheat bran is 7.5g/g. This indicates the improved water absorption of the fibre after cell wall components have been partly removed. Other commercial dietary fibres extracted from wheat straw and sugar beet have WHC of 6.3 and 7.9g/g, respectively.

EXAMPLE 9

10 The protein fraction, which contains substantial amounts of aleurone proteins, produced as described in Example 7, have a very interesting chemical composition, functionality and is an ideal raw material for further processing.

A typical composition of the protein fraction is: dry matter 98%, protein 40%, sugar 3%, fat
 15 18%, non-sugar carbohydrates 32% and minerals 5%.

In order to determine the effect of protease treatment on the functionalities of the protein fraction, a protein sample was subjected to a mild protease treatment and the samples analysed for dry matter and protein solubility, emulsifying capacity and emulsifying
 20 stability.

The results are shown in Table 2, and clearly indicate the possibilities to further improve some important functionalities of the protein fraction.

25 TABLE 2

Parameters analysed	Protein fraction	Protein fraction treated with protease
Dry matter solubility (%)	19.7	38.1
Protein solubility (%)	18.4	55.5
Emulsifying capacity (%)	52.5	90.6
Emulsifying stability (%)	47.5	86.0

Ink. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassan

15

EXAMPLE 5

Amongst the various end-uses of the germ-rich fraction of EXAMPLES 1-4 one could describe meat products such as hamburgers, sausages and meatballs. In such end-uses germ-rich fraction could replace meat, soy protein concentrate and isolate, but also milk casein and caseinates, to mention just a few. It is therefore important to test the overall performance of the germ-rich fraction with regards to emulsifying and binding capacity, taste, etc.

A trial set up to test the feasibility of incorporating various germ-rich fractions extracted from wheat bran into a traditional meat ball recipe consisted of meat, garlic, premix and water.

The following spray dried fractions were tested:

- Germ-rich fraction extracted from short milling wheat bran - (I)
- Germ-rich fraction extracted from conventional milling wheat bran - (II)
- 1:1 mix of whey and II, on dry matter basis - (III)

Meatball recipes were tested without germ-rich fraction (control recipe) or with 2.5% inclusion of samples I, II or III. Meatballs were analysed for weight loss, taste, texture and colour after frying.

The results are described in the table 3 below.

TABLE 3

Recipe tested	Weight loss after frying (%)	Colour	Texture
Control (meat, garlic, premix and water)	23.4	Reference	Reference
Control + 2.5% of I	21.3	Slightly darker	Slightly tougher
Control + 2.5% of II	20.8	Similar	Similar
Control + 2.5% of III	18.0	Similar	Slightly more tender

The overall conclusion was that the samples performed well as additives in a meat ball recipe, and were particularly interesting as they all decreased the weight loss after frying.

30

16

Ink. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassen

END-USES

Germ-rich fraction

5 The high protein content of the germ-rich fractions makes it an ideal substitute for existing expensive proteins from animal and vegetable origin. Additionally, the germ-rich fraction because of the nature of its protein, the presence of high quality oil and phospholipids also exhibit interesting functionalities such as emulsification, texture and binding.

10 One can list, as examples, the following existing products, which can be replaced by the germ-rich fraction in the food industry:

Animal protein: casein and caseinates, plasma protein and egg white

Vegetable proteins: soy protein concentrates and isolates, texturized soy, hydrolysed gluten and potato protein,

15

Generally, the above products can be used as meat extenders and texturizer ingredients in hamburger, sausage, and meat balls production to mention a few. Or, as a casein replacer in the production of sausage, spreads, dressings, etc.

20

In the feed industry, the germ-rich fraction is an ideal ingredient for high value feeds such as calf milk replacer, starter feeds for calves, piglets and chicks, fish feeds and pet food. In such applications it can substantially replace the use of soy proteins (texturized soy, concentrate and isolate), potato protein, hydrolysed gluten, high quality fish meal, plasma protein, and dry milk products such whey protein concentrate, whey and skimmed milk.

25

Endosperm-rich fraction

The endosperm-rich fraction can be used in the food industry primarily as a gluten replacer, but also to partly substitute soy proteins in the production of various meat and vegetable products. In feed applications, it can partly replace gluten, soy and milk proteins as an ingredient to calf milk replacer, piglet started feed and fish feed.

30

Ink. t. Patent- och reg.verket

2001-10- U 4

Huvudfaxen Kassen

17

Fibre fraction

The fibre fraction consisted of pericarp and aleurone can be used as an interesting source of gluten- and phytate-free fibre source to replace conventional cereal brans. The main end-uses as food would be in the baking industry and breakfast cereal.

5

One preferred embodiment of a plant for carrying out the invention is shown in the attached drawing,

wherein FIG. 1 a set-up for carrying out the process of the invention;

FIG. 2 shows a set-up for carrying out a preferred embodiment of the invention; and

10 FIG. 3 shows the fractionation of the bran in an over-view.

In FIG. 1, 1 denotes a suspension and hydrolysis vessel 1 and connected to a wet mill 2 which by its milling action increases the active surface of the hydrolysis. The slurry is allowed to pass the wet mill 2, 1 to 3 times. The slurry is then transferred to a 2-

15 phase decanter 6 via a heat exchanger 3, which decanter 6 separates the bran and liquid (water and protein) phases. The bran having a dry matter content of about 40%, is then further washed once using water in a suspension vessel 4, and is allowed to pass a second decanter 6, again. Then a pure bran obtained has a dry matter content of 95% after drying in a ring drier 10. The collected liquid phases, having a dry matter content of 5%, are

20 transferred to a separator 7, via a mixing tank 5, in which separator 7 insoluble protein is separated off. The liquid phase from the separator 7, having a dry matter content of 2%, and comprising water and soluble proteins is allowed to pass an ultra filter 8 having a molecular cut of 50 kD. Depending on different requirements this cut can vary between 20 and 100 kD. After the ultra filter 8 a liquid phase is obtained which is evaporated in an evaporator 9, and concentrated to a syrup, having the dry matter content of 75%. The concentrate of the

25 ultra filtration is either isolated as such, or added to the fraction of insoluble proteins, which latter has a dry matter content of 9.5%. The protein-sugar fraction obtained is spray-dried in a spray drier 11 to provide a concentrated, dried protein-sugar fraction, and the protein-oil fraction is dried in a spray-drier 11 to provide a concentrated, dried protein-oil fraction.

30

One preferred embodiment of a plant for carrying out the invention related to clean bran separation is shown in the attached drawing, FIG. 2 wherein 1 denotes a suspension and hydrolysis vessel to which a wet mill 2 is connected. The reaction mixture is intermittently

Ink. t. Patent- och reg.verket

2001-10-04

Muvudfaxen Kassen

18

pumped through the wet mill 2 (from 1 to 3 times). The hydrolyzate is then inactivated in a heat exchange 3 and transferred to a two-phase decanter 4, which decanter 4 separates the insoluble (insoluble dietary fibre) from the soluble phase. The insoluble phase having a dry matter content of about 35% is dried to approximately 95% dry matter in a ring drier 5. The soluble phase, having a dry matter content of approximately 3%, is pumped through another two-phase decanter 7, via a holding tank 6, in which two-phase decanter 7 protein-rich fraction is separated off. The protein-rich fraction is then dried to about 95% dry matter in spray drier 8. The soluble (liquid) phase from the two-phase decanter 7, having a dry matter content of approximately 3%, and comprising water and primarily soluble hemicellulose, oligosaccharides and sugar is allowed to pass an ultra filter 9 having a molecular cut of 20 kD. Depending on different requirements this cut can vary between 10 and 50 kD. The retentate (fraction retained in the ultrafilter) from ultrafilter 9 is evaporated to a syrup concentration of at least 75% solids in an evaporator 10 or alternatively in a spray drier such as 8. And, the permeate (fraction not retained in the ultrafilter) from ultrafilter 9 is preferably evaporated to a syrup concentration of at least 75% solids in an evaporator 11.

Applications

Insoluble dietary Fibre

This is the remaining fibre after the soluble components and a proportion of hemicellulose has been removed from rye bran. The insoluble dietary fibre is a cleaned cereal fibre containing low levels of phytic acid. Because the fibre has already been partially "digested" enzymatically, many beneficial compounds derived from the cell wall are available to the gut for absorption.

25

The insoluble dietary fibre has high water binding capacity, i.e. typically 100% higher than that of wheat bran. This provides increased gut transit (digesta flow). The remaining pentosans are more accessible to the gut wall (cholesterol reducing) due to the fractionation process. Because of the increased availability of lignin type materials and other antioxidants within the fibre, various health benefits can be claimed. Specifically the lignans and polyphenolics are known to mimic estrogens (female hormones), and more recently have been found to help preventing various types of cancers. This has been verified for rye products.

30

Ink. t. Patent- och reg.verket

2001-10-04

Muvudlexen Kassar

19

5 The insoluble dietary fibre is very useful as a food additive as a water binding and texturising ingredient and is good for the gut as a dietary supplement. Specifically the high water binding capacity and beneficial effect on bowel function makes it an interesting product for the biomedical market.

10 Additionally the insoluble dietary fibre is also a good raw material for the further extraction (enzymatically) of lignins, ferulic acids, lignans etc. which are natural antioxidants and potential anticancer agents. These can be used in many biomedical and "cosmetic pharmaceutical" applications such as lotions, creams and moisturizers. The ferulic acid is an effective UV absorber and as such can be used in a sunscreen.

15 Insoluble dietary fibres are rich in accessible lignans and residual pentosans / hemicelluloses. Bacteria present in the colon convert plant lignans to mammalian lignan, enterolactone, using hemicelluloses as a fermentation medium. These compounds mimic oestrogens and appear to have a tangible, demonstrable effect on the suppression of hormone related cancers, eg breast, ovarian and prostate cancers. Rye insoluble dietary fibre specifically contains the lignans Secoisolariciresinol (SECO) and matairesinol (MAT) which are known precursors of enterolactones. The insoluble dietary fibre from wheat also
 20 contains these lignans, but the effect is not demonstrated in wheat. It is important to state that in this fraction the lignans are supplied in an accessible form, as the cell wall is already partially enzymatically digested, along with their natural synergistic partners, the arabinoxylan hemicelluloses remaining on the fibre.

25 A further health claim is the improved water holding capacity of the fibre as compared to normal bran. This increases digesta flow and reduces gut transit times, with concomitant benefits to health.

Aleurone Rich Protein

30 This is the protein that is derived from the aleurone cell layer and is both a functional and nutritionally valuable material, rich in essential amino acids.

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassar

20

In the food business it would be used as a emulsifier, foam stabilizer and texturiser. In addition there is high potential as a protein supplement.

5 Soluble Hemicellulose

This is the major cell wall non-cellulose polysaccharide in rye bran. It can be produced with high molecular weight and high solubility (the combination of these two properties is the powerful aspect). Because the product is a pentosan (arabino-xylan) it is low calorie and beneficial for gut health. The product can be produced with or without ferulate side chains or free ferulic acid and other antioxidants and is a free flowing cream powder.

Due to its composition and high water binding capacity it is ideal to use as a thickener, gellant, stabilizer, soluble dietary fibre and fat replacer. As a thickener and gellant it is interesting in the food industry as an additive in soups, margarine, deserts, pâtés, sauces etc. As a stabilizer it is a cheaper alternative for modified starch (made from wheat, maize etc), modified cellulose, gums (guar gum and carrageenan gum), alginates (seaweed), gelatin (cheap but problem with BSE) and pectin (fruit peel & sugar beet). Finally, it has a big potential in drinks because it is an excellent source of soluble dietary fibre alongside its stabilizing properties.

20

It is possible to supply the pentosan with ferulate side chains, and in this form the substance will gel in combination with oxygen and enzyme. As such it is an interesting material, for example, for wound dressings as it will keep the skin in a hydrated state and therapeutic agents can be added.

25

From rye and wheat, this is almost exclusively arabino-xylan (pentosan) hemicellulose. This is readily fermented in the colon, is low calorie to the human and is reported to generate butyrate as a short chain fatty acid (SCFA) end product after fermentation. This is the most "healthy" SCFA according to recent studies as it is a preferred source of energy to epithelial cells lining the colon. The health benefits of adding an enriched, available source of arabinoxylan to the diet may therefore be far-reaching.

30

Ink. t. Patent- och reg.verket

2001-10-04

Muvudfaxen Kassan

21

The fraction is a perfect soluble dietary fibre, with all of the concomitant health benefits. Arabinoxylans are also thought to be excellent binding sites for secondary bile acids, as a consequence of the rigidity of parts of the molecular chain and the occurrence of relatively hydrophobic domains on the polymer. This is thought to reduce any potential carcinogenic effects.

In addition, the fraction contains ferulic ester side chains to a proportion of the polymer, with concomitant free radical stabilising and anti-oxidant properties.

It is important to emphasise that the arabinoxylan concentrated in this fraction is not normally available to the gut and colon if presented as a part of a normal diet or even from conventional bran.

In oat, this fraction is rich in β -glucan with all of the documented beneficial effects of this polysaccharide. There is a tangible benefit in supplying a concentrated β -glucan of this nature as the normal ingestion of oats does not supply sufficient material for the full effects to be realised. Purified β -glucan can be purchased but is very expensive because of the extensive purification regime. It is important to realise that this high purification is required to remove the chemicals utilised in the extraction process. It is suspected that natural synergistic partner compounds are removed in such a process, whilst these materials should still be present in the present process fraction.

Oligosaccharide Syrup

This is derived from the hemicellulose fraction and is a 100% soluble dietary fibre of low molecular weight and low viscosity. The oligosaccharide syrup can be produced with lignans, ferulic acid and other antioxidants and is extremely soluble and hygroscopic.

It has a high potential in the drinks industry as it has low viscosity, is a good source of dietary fibre and gives good mouth feel and texture.

In combination with glucose syrup it could be used as a sweetener and energy source for drinks, cereal bars etc. As it is rich in ferulic acid, pentosans and solubilised lignans, one

Ink. t. Patent- och reg.verkst

2001-10-04

Huvudfaxen Kassar

22

can also claim the related health benefits. It is very important to supply lignans in the presence of pentosan oligomers if the full cancer prevention effect is to be expressed and realised: precisely the situation in this fraction.

5

The combination of glucose syrup and oligosaccharide syrup is also ideal in applications where one requires increased dietary fibre content and increased water binding capacity without thickening.

10 The oligosaccharide syrup is the low molecular weight fraction of the solubilised arabinoxylans along with other low molecular weight components solubilised from the cell wall. This includes dissolved lignin fragments, phenolic compounds such as ferulic acid and lignans. As with the insoluble dietary fibre, the presence of lignans with arabinoxylan gives rise to claims for cancer preventative roles for rye and wheat derived fractions. In this case,
 15 the arabinoxylans are present as oligomers and the lignans are very available in the syrup with a potential high accessibility for the gut. This should increase the rate of conversion of the plant lignans to enterolactone with a potentially larger impact on cancer prevention.

Furthermore, the presence of high concentrations of oligomeric arabinoxylan provides a
 20 ready fermentation substrate for the production of beneficial SCFAs such as butyrate, with benefits as described for the hemicellulose fraction.

This fraction, especially in the case of rye, is probably the most interesting in the present context being an excellent source of arabinoxylans, lignans and phenolic antioxidants in
 25 very accessible forms along with relevant synergistic partner compounds.

Germ Rich Protein

The germ rich protein is one of the products that are produced from the soluble fraction. It is one of the most valuable products because it contains the rye germ oil. Typically, it contains
 30 55% protein and 17-20% oil, the latter being rich in poly-unsaturated fatty acids and in beneficial materials for cholesterol control.

Link t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kossan

23

The germ rich protein is a good functional protein and can also substitute soy protein in many applications. Its good texturising effects are very interesting for the meat industry (meatballs, hamburgers, pate etc.). Properties such as solubility and emulsification can be further improved by enzymatic hydrolysis.

The germ rich protein is of great interest as a functional food ingredient, especially in the case of rye, primarily because it contains the germ oil (see below).

10 Cereal Germ Oil

All of the germ oils isolated using the present process are interesting from a health point of view. All contain the vitamin E tocopherol complex, along with tocotrienols, vitamin B and polyunsaturated fatty acids. The case of rye germ oil is especially significant, as it is one of nature's richest sources of β -sitosterol, a documented cholesterol inhibitor, alongside tocotrienol, a cholesterol burner. Furthermore, the form of the sterol in the oil is suitable for adsorption. The present process represents a commercial route to rye germ oil. A formulation in margarines and spreads is suggested as a first step to commercial utilisation.

Rye Germ Oil

The germ oil is derived from the germ rich protein and is a very high quality food grade oil and ingredient. It is extracted without using any solvent, and it contains no preservatives or additives. It is a good source of poly and mono unsaturated fat, has a good flavour, is rich in vitamin E and can be suspended easily. As a flavouring component it is good in wheat or rye based products (cereals, baking goods, biscuits etc), deserts, ice creams etc. It can also be useful as an ingredient in fat and oil formulations, juices etc. with natural vitamin E.

Rye germ oil is particularly rich in naturally occurring β -sitasterol, a cholesterol lowering compound and tocotrienol, a cholesterol "burner". These materials can be classified as "natural synergistic partners", an important factor in the functional food area. This massively increases the potential of the oil as a value-added nutraceutical ingredient in foods such as margarines and spreads.

Int. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassa

24

Rye germ oil is also a good UV blocker and therefore together with ferulic acid it can be ideal as a component of a sun tan lotion. Its emulsifying properties make it very suitable as an emulsion stabilizer and as an emollient ingredient for skin creams. It should be mentioned that there is presently no reasonable scale commercial production of rye germ oil and the present process is the only realistic way to achieve this.

Endosperm Rich Protein

This protein fraction is mainly derived from the residual endosperm proteins in rye bran. It contains 35-40% protein, much of which is highly soluble. It is also particularly rich in pentosans (soluble dietary fibre), has a high water holding capacity and has a light colour.

The emulsifying, water binding and foam stabilising properties are equivalent or better than those of other commercial proteins like caseinates, soy protein concentrate and modified wheat gluten. The endosperm rich protein is very suitable to be used as an ingredient in milk replacer formulae (both for humans and calves), sauces, mayonnaise, dressings etc.

Because it contains high amounts of pentosans and associated ferulic acid there are extra health and functional benefits. In the cosmetic industry the stabilising, emulsifying and water holding properties are ideal.

A combination of the endosperm rich protein and soluble hemicellulose is interesting in a number of food and biomedical applications, because of the emulsifying effect of the endosperm rich protein and the soluble pentosans and the thickening effects of the soluble hemicellulose.

The endosperm rich oil is rich in pentosan hemicelluloses, mainly arabinoxylans in rye and wheat, or β -glucans in the case of oat. The claimed health benefits are therefore as described already for these fractions.

Glucose Syrup

This is the glucose produced from enzymatic degradation of residual starch of the bran and is a more pure product compared to molasses. As such it is ideal as a feedstock for industrial

5 fermentation since it produces less waste products. Ethanol and citric acid industries are therefore ideal consumers of very large quantities of such a product. The production of single-cell protein for the feed and food markets can also be considered. Alternatively, it can be used as a sweetening syrup in the standard fashion.

Defatted Germ Rich Protein

10 This is the protein that remains after the rye germ oil has been extracted from the germ rich fraction and has at least 70% protein content (is in the same range as soy protein concentrates). It is also a good functional protein, has an extremely high fat binding capacity and can be easily upgraded enzymatically to increase solubility, emulsion and foam stabilisation properties.

15 The product is an excellent stabiliser for water in oil emulsions and is interesting as a meat texturiser or extender in sausages, burgers, patées etc. The defatted germ rich protein is a functional protein that can easily replace soya proteins and contains phospholipids, natural lecithins and glycolipids.

20 It has a high potential in cosmetic formulation as an emulsion stabiliser because it contains natural lecithins.

CLAIMS

1. Process for the wet fractionation of cereal bran components,
 characterized in
 that bran is first subjected to a combination of enzymatic treatment with enzymes of the
 5 group starch- and phytate-hydrolysing enzymes, and wet milling, followed by an optional
 step of enzyme inactivation by wet heat treatment, and the next step whereby the insoluble
 phase containing both pericarp and aleurone fractions are separated by centrifugal forces
 from the aqueous phase containing germ and residual endosperm components to produce a
 clean bran, and that the aqueous phase is further separated by centrifugal force into a germ-
 10 rich fraction and an endosperm-rich fraction, and that the proteins contained in the
 endosperm-rich fraction are concentrated.
2. Process according to claim 1,
 wherein cereal brans are the fibrous-residue resulting from a primary grain milling, i.e. after
 15 the separation of the endosperm fraction, of wheat, rice, barley, oat, rye and triticale, and
 having variable chemical compositions, presence of anti-nutritive factors, and presence of
 various anatomical fractions, i.e. pericarp, germ, and residual endosperm.
3. Process according to claim 1,
 20 wherein the combination of wet milling with enzymatic treatment is arranged to increase
 substrate accessibility thereby improving the overall hydrolysis performance and the
 subsequent separation of both insoluble and soluble fractions of varied density/solubility.
4. Process according to claim 1,
 25 wherein the enzymatic treatment is accomplished by using a non-starch degrading enzyme
 in the form of a polysaccharidase of amylases and/or amyloglucosidases.
5. Process according to claims 1-4,
 wherein a further enzymatic treatment is carried out using at least one non-starch degradable
 30 polysaccharidase in the form of cellulases, hemicellulases mainly xylanases, beta
 glucanases, and pectinases, and/or phytases.

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassar

27

6. Process for the wet fractionation of cereal bran substantially free of soluble compounds according to claim 1,
 wherein such clean bran is subjected to a combination of enzymatic treatment with specific
 5 enzymes of the group xylanase under strictly controlled hydrolysis conditions, and
 intermittent wet milling, followed by an optional step of enzyme inactivation by wet heat
 treatment.
7. Process according to claim 6,
 10 wherein the inactivated hydrolysate is then fractionated by centrifugal forces into an
 insoluble phase containing primarily cellulose, lignin, less accessible hemicellulose, residual
 aleurone cells and cell wall bound proteins, and an aqueous phase containing soluble
 hemicellulose, oligosaccharides, sugars and proteins, and that the aqueous phase is further
 separated by centrifugal force into protein-rich fraction and a carbohydrate-rich fraction,
 15 and that the carbohydrate-rich fraction is further separated by size exclusion technique into a
 hemicellulose-rich fraction (large molecular size) and an oligosaccharide-rich fraction
 (small molecular size).
8. Process according to claims 6-7,
 20 wherein cereal bran substantially free of soluble compounds, both in water and/or non-polar
 solvents are derived from wheat, rice, barley, oat, rye or triticale.
9. Process according to claims 6-8,
 wherein the combination of intermittent wet milling with enzymatic treatment is arranged to
 25 increase substrate accessibility to the cell wall degrading enzymes thereby improving the
 overall hydrolysis performance and the subsequent separation of the various fractions by
 density/solubility and molecular size.
10. Process according to claims 6-9,
 30 wherein the enzymatic treatment is carried out using at least one non-starch degradable
 polysaccharidase in the form of cellulases, hemicellulases mainly xylanases, beta
 glucanases, and pectinases, and/or phytases.

Ink. t. Patent- och reg.verket

7001-10-04

28

Huvudfaxen Kassar

11. Process according to claim 10,

wherein the enzymatic treatment is accomplished by using xylanases with high beta 1-4-xylanase (pentosanase) activity.

5 12. Protein fraction derived substantially from the germ and produced according to claims 1-4,

wherein the said fraction contains at least 40% protein and 10% oil on dry matter basis and exhibits an increased shelf life with regards to resistance to oxidation compared to the original bran, and that the said fraction contains less than 1% fibre, and it retains the
 10 emulsifying capacity of wheat germ.

13. Protein fraction according to claim 12,

wherein liquid whey is incorporated in to the said fraction at levels varying from 20 to 80% by weight on dry matter basis, and that the final mixture is dried.

15

14. Protein fraction derived substantially from the residual endosperm and produced according to claims 1-4,

wherein the said fraction contains at least 35% protein and 10% sugar and less than 2% oil and 1% fibre.

20

15. Protein fraction according to claim 14,

wherein liquid whey is incorporated in to the said fraction at levels varying from 20 to 80% by weight on dry matter basis, and that the final mixture is dried.

25 16. Fibre fraction produced according to claims 1-4,

wherein the said fraction consists of cell wall components of bran (>85%) and aleurone proteins (>10%), and substantially free of gluten and starch.

17. Sugar fraction produced according to claims 1-4,

30 wherein the said fraction is originated primarily from the residual endosperm and it contains more than 65% sugars (such as glucose, maltose and malto-triose) on dry matter basis.

18. Insoluble dietary fibre according to claim 16, used for recovery of lignans.

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassan

29

19. Insoluble dietary fibre according to claims 6-7, used for recovery of lignans
20. Aleurone rich protein produced in accordance with claims 6-11.
- 5 21. Soluble hemicellulose produced in accordance with claim 6-11.
22. Oligosaccharide syrup produced in accordance with claim 6-11.
- 10 23. Germ oil produced in accordance with claims 1-5.
24. Rye germ oil produced in accordance with claims 1-5.
25. Defatted germ rich protein produced in accordance with claim 1-5.
- 15 26. Use of a protein fraction, as described in claims 10-13, and 17, in feed and food applications to replace other protein products from vegetable and animal sources.
- 20 27. Use of a fibre fraction, as described in claim 16, 18, 19, in feed and food applications to replace other insoluble fibrous products or as a raw material for further processing.
- 25 28. Protein fraction derived substantially from the aleurone cells and produced according to claims 6-11,
wherein the said fraction contains at least 40% protein and 15% oil, less than 1% insoluble fibre on dry matter basis, substantially free of gluten and starch and with a high emulsifying capacity.
- 30 29. Protein fraction according to claim 28,
wherein the fat can be optionally removed by conventional organic solvent extraction or preferably by supercritical carbon dioxide extraction to yield a defatted protein and an oil fraction.
30. Protein fraction according to any of claims 10, 12, 28, and 29,

Ink. t. Patent- och reg.verket

2001-10-04

30

Huvudfaxen Kasse

wherein proteases are incorporated in to the said fraction in wet state and at controlled temperature and pH conditions, and the resulting protein hydrolysate has enhanced functionalities such as solubility, emulsifying and foaming capacities.

5

31. Insoluble fibre fraction produced according to claims 6-11, wherein the said fraction consists primarily of cell wall components with a relative lower hemicellulose content compared to the original cleaned cereal bran, substantially free of gluten and starch (<1% on dry matter basis) and with a high water holding capacity (>8g water/g dry product).

10

32. Soluble hemicellulose fraction produced according to claims 6-11, wherein the said fraction consists primarily of high molecular weight hemicellulose of preferably above 20 kDa (>30%), which also contains proteins (<10%) and monosaccharides (<10%), and is substantially free of gluten and starch (<1% on dry matter basis).

15

33. Soluble oligosaccharide fraction produced according to claims 5-9, wherein the said fraction consists primarily of low molecular weight hemicellulose sub-units of below about 20 kDa (>30%), which also contains proteins (<10%) and monosaccharides (<20%), and is substantially free of gluten and starch (<1% on dry matter basis).

20

34. Use of a protein fraction, as described in claim 29-30, in feed and food applications to replace conventional protein ingredients and functional proteins both from vegetable and animal sources.

25

35. Use of an insoluble fibre fraction, as described in claim 31, in feed and food applications as a water binder and texturizer to replace other insoluble fibres, or as a raw material for further processing to extract the remaining cellulose and lignin.

30

36. Use of a soluble hemicellulose, as described in claim 32, in feed, food and chemical applications as a gellant, thickener, emulsifier and/or as a source of soluble dietary fibre, or as a raw material for further processing to obtain other functional hemicelluloses.

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfoxen Kassan

31

37. Use of a soluble oligosaccharide, as described in claim 33, in feed and food applications as a functional soluble dietary fibre or low calorie sweetener.

5 38. Set up for carrying out the process according to claims 1-4, characterized in

that it comprises a hydrolysis vessel (1), a wet mill (2), a heat exchange for enzymatic inactivation (3), washing tanks (4), a mixing tank (5), a decanter (6), a separator (7), an ultra-filter (8), an evaporator (9), and optionally dryers (10).

10

39. Set up for carrying out the process according to claims 6-11, characterized in

that it comprises a hydrolysis vessel (1), a wet mill (2), a heat exchange for enzymatic inactivation (3), decanters (4 and 7), dryers (5 and 8), a holding tank (6), an ultra-filter (9) and evaporators (10 and 11).

15

40. Process according to claims 1-5,

wherein the enzymatic treatment is carried out at a pH of 4 to 7.5 and at a temperature of from 50 to 90°C, at an enzymatic activity of 200 to 1500 IU/g of substrate.

20

41. Process according to claims 6-11,

wherein the enzymatic treatment is carried out at a pH of 4 to 7, preferably 4.5-5.5, and at a temperature of from 35 to 80°C, at an enzymatic activity of 200 to 1500 IU/g of substrate.

25

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfoxen Kassan

32

ABSTRACT

A process for the fractionation of cereal brans substantially free of soluble compounds is described. In particular, this invention describes a two step process, in which the said cleaned bran is subjected to a combination of enzymatic treatment with xylanases and wet milling, and a second step consisting of sequential centrifugation and ultrafiltration, which aims at physically separating the main fractions, i.e. insoluble phase (remaining cell wall components), protein-rich fraction, soluble hemicellulose and oligosaccharide. This invention also describes a process that maximizes the extraction rate of valuable cell wall components and aleurone cells from previously cleaned bran.

10 (FIG 1)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146

Ink. t. Patent- och reg.verket

7U01-10- 0 4

Huvudfoxen Kassan

FIG. 1



Ink. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassa




FIG. 2

Ink. t. Patent- och reg.verket

2001-10-04

Huvudfaxen Kassar




FIG. 3

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.